

EFFICACY OF AQUEOUS SEED EXTRACTS AGAINST *PECTOBACTERIUM CAROTOVORUM* CAUSING BLACK LEG AND SOFT ROT OF POTATO

G. Biswal* and N. K. Dhal

Department of Plant Pathology, College of Agriculture, Odisha University of Agriculture & Technology.

Received on: 29/08/2020

Revised on: 19/09/2020

Accepted on: 09/09/2020

*Corresponding Author

G. Biswal

Department of Plant
Pathology, College of
Agriculture, Odisha
University of Agriculture &
Technology.

ABSTRACT

Studies were conducted in the Department of Plant Pathology, Odisha University of Agriculture and Technology, Bhubaneswar following inhibition zone technique to test the bio-efficacy of seed extracts *in vitro* against *Pectobacterium carotovorum* causing pre-emergence and post-emergence rotting, black leg in field soft rot in storage in potato. The seeds used in the tests were *Terminalia chhebula* (Chhebolic myrobalan), *T. belerica* (Beleric myrobalan), *Embllica officinalis* (Indian goose berry), (*Azadirachata indica* (Neem), (Greater cardamom), *Rauvolfia serpentina* (Snake root), *Coriandrum sativum* (Dhania), *Cuminum cyminum* (Cumin), *Nigella sativa* (Black cumin), *Foeniculum vulgare* (Fennel), *Piper nigrum* (Black pepper), *Cassia fistula* (Indian laburn), *Cassia tora* (Senna tora). The results revealed that *R.serpentina* exhibited maximum size of inhibition zone 13.53 followed by *A.subulatum*(11.33) against the test bacterial species. The inhibition zone was found to be 10.0mm in seed extracts of *T.chebula* while 9.30mm in *F.vulgare* 8.68 in *C.cyminum* and 8.03mm in *N.sativa*. There was no significant difference in antibacterial activity between *A aromaticum* (7.97) and *A.indica*(7.92). The antibacterial activity was same (7.01mm) in *T.bellirica* and *C.viscosa*. In *P.nigrum*, *C.fistula* and *C.tora* expressed same inhibiting activity (6.95mm). Minimum inhibition zone was observed in *E.officinallis* (6.68mm).

KEYWORDS: Aqueous plant extracts, Antibacterial activity, black leg and soft rot.

INTRODUCTION

Potato (*Solanum tuberosum*) is the most potential and nutritionally superior crop for fighting against hunger in both developing and under developed countries. The world wide occurrence of potato diseases mainly revealed the presence of 30 fungal, 7 bacterial and 36 viral diseases causing loss individually or collectively to the crop. The blackleg and soft rot caused by *Pectobacterium carotovorum* is an important disease of potato of the country, damaging the crop during pre and post emergence stage growing period harvest and post harvest operations like storage, transport and also at consumer level. The disease was found to be severe in immaturely harvested and peeled tubers while transported under poorly ventilated condition and under normal package of practices, i.e. 4% in north western plains and 10% in eastern and peninsular India and 4-8% in hills (Somani and Shekhawat, 1990). In different locations of Odisha the black leg incidence varied from 1.35 to 4.36% in growing period and 4.19 to 6.47% of soft rot of tubers during harvest (Biswal and Dhal, 2013). The use of plant products have remarkable effects in plant disease management (Cowan, 1990 and Newman *et al.*, 2000). The use of organic farming is now gaining popularity. Different parts of many plants have antimicrobial properties. In this context the seeds of some commonly grown trees, weeds, medicinal plants

and spices were tested against *P. carotovorum* causing black leg and soft rot of potato. The seeds used in the studies were *Terminalia chhebula* (Chhebolic myrobalan), *T. belerica* (Beleric myrobalan), *Embllica officinalis* (Indian goose berry), (*Azadirachata indica* (Neem), (Greater cardamom), *Rauvolfia serpentina* (Snake root), *Coriandrum sativum* (Dhania), *Cuminum cyminum* (Cumin), *Nigella sativa* (Black cumin), *Foeniculum vulgare* (Fennel), *Piper nigrum* (Black pepper), *Cassia fistula* (Indian laburn), *Cassia tora* (Senna tora). Hence the present investigation comprising *in vitro* testing of abovementioned seed extracts was conducted in the Department of Plant Pathology, Odisha University of Agriculture and Technology, Bhubaneswar following inhibition zone technique (Valgas *et al.*, 2007). The study was undertaken keeping in view of use such extracts as tuber treatment at planting time, basal drenching in field and also tuber treatment in storage after harvest.

MATERIALS AND METHODS

The healthy seeds of fifteen plants, i.e. *Terminalia chhebula* (Chhebolic myrobalan), *T. belerica* (Beleric myrobalan), *Embllica officinalis* (Indian goose berry), (*Azadirachata indica* (Neem), (Greater cardamom), *Rauvolfia serpentina* (Snake root), *Coriandrum sativum* (Dhania), *Cuminum cyminum* (Cumin), *Nigella sativa*

(Black cumin), *Foeniculum vulgare* (Fennel), *Piper nigrum* (Black pepper), *Cassia fistula* (Indian laburn), *Cassia tora* (Senna tora) were collected (Table 1). These were washed several times in sterilized water and air dried. Fifty grams from selected seeds along with 50ml of double distilled water were taken grinded with the help of pestal and mortar to a fine pulp. The pulp was filtered through two layers of muslin cloth and gently pressed to get maximum filterate. The filterate from each plant part was collected and kept separately in different sterile specimen tubes and centrifused at 1500 rpm for 15minutes. The supernatant liquid was drawn carefully into a 5ml syringe and then passed through membrane filter of 0.45nm size to sterilize the extract. The filter sterilized extract of each part collected in sterilized specimen tube with screw cap and stored in deep freeze maintain edat -20° C. The extracts were evaluated *in vitro* following the inhibition zone technique. In this technique, two drops of bacterial suspension of each test

bacterium was transferred on to the petriplate containing NSA medium and spreaded over the surface of the medium with the helpof a sterilized glass spreader. Three sets of Hi-media discs(5mm), soaked for one minute in each plant extracts were placed on the media surface of each petriplate at the equidistance from the centre. In each set four numbers of discs were used to hold sufficient quantity of the plant extract. Two sets of petridishes were used for testing each plant extract petriplates were incubated at $27\pm 1^{\circ}$ C for 24 hours in a BOD incubator. After the incubation period, the petriplates were examined for development of inhibition zone around the discs. The diameters of each zone of inhibition was measured and recorded and analysed stastically to assess the antimicrobial properties of plant extracts against each test bacterium (Gomez and Gomez,1984). In control the paper discs were soaked in sterilized water.

Table 1: Scientific name, common name, family and traditional uses of test plants.

Sl. No.	Scientific Name	Common name (English)	Family	Traditional uses
1	<i>Terminalia chebula</i>	Chebulic myrobalan	Combretaceae	It is an important ingradient for Ayurvedic medicine Triphala used intreatment of constipations, colic pain, kidney dysfunction, eye diseases and sore throat,(https://vikaspedia.in/agriculture , Basa <i>et al</i> ,2017)
2	<i>T.belerica</i>	Belericic myrobalan	Combretaceae	It is also an ingradient for Ayurvedic medicine Triphala used in treatment of constipations, colic pain, kidney dysfunction, eye diseases and sore throat. Seeds are edible (Kumar <i>et al</i> , 2018)
3	<i>Emblica officinalis</i>	Aonla	Euphorbiaceae	ingradient for Ayurvedic medicine Triphala used in treatment of constipations, colic pain, kidney dysfunction, eye diseases and sore throat (Sharma <i>et al</i> ,2003)
4	<i>Amomum subulatum</i>	Greater cardamom	Zingiberaceae	It is analgesic,antimicrobial cardiac stimulant,carminative,diuretic stomachi (Bisht <i>et al</i> ,2011)
5	<i>Rauvolfia serpentina</i>	Snake root	Apocynaceae	Used for various aliments such as snakebites, insomnia, hypertension and insanity (Singh <i>et al</i> , 2017. Negi <i>et al</i> , 2014)
6	<i>Cleome viscosa</i>	Wild mustard	Cleomaceae	Used against fever, diarrhea, cardiac stimulant and carminative (Perumal Samy <i>et al</i> 1999)
7	<i>Coriandrum satium</i>	Coriander	Apiaceae	It is ntidiabetic, anti-inflammatory and loers cholesterol. It is used as diuretic, carminative, stimulant, nagelsteic, antihelmtic, hypoglycaemic (Waheed <i>et al</i> ,2006)
8	<i>Cuminum cyminum</i>	Cumin	Apiaceae	Seeds Food additive, popular spice, flavouring agent in many cuisines. It is used against hypolipidemica, cancer and diabetes (Mnif and Aifa, 2015)
9	<i>Azadirachata indica</i>	Neem	Meliaceae	Anti oxidant, antimalarial, antimutagenic,anticarcinogenic,anti inflammatory, antihyperglycaemic, antiulcer and antidiabetic purposes(Venugopalan and Visweswaran,2013)
10	<i>Cassia tora</i>	Senna tora	Fabaceae	Used against leprosy, bronchitis and cardiac disorders(Maity <i>et al</i> ,1998)
11	<i>Nigella sativa</i>	Black cumin	Ranunrulaceae	Commonly used for culinary and medicinal purposes as a remedy of hypertension and diabetes and as hypoglycemic, anti-inflammatory, antiulcer and broncho dilator(Bereksi <i>et al</i> ,2018)
12	<i>Foeniculum vulgare</i>	Fennel	Apiaceae	Fennel seeds helps in digestion, prevents acne, mouth freshner, beats bad breath(Al-Timimi ,2019)
13	<i>Piper nigrum</i>	Black pepper	Piperaceae	It is used as spice. It exhibit sedating, detoxification, hypotensive and anticancer activities. Butt <i>et al</i> 2012
14	<i>Amomum aromaticum</i>	Aromatic cardamom	Zingiberaceae	Seeds are used to make a gangle or mouth wash to treat toothache, gingivitis and paradontosis. Seeds are antibacterial and use against

				stomachic, alleviate dyspepsia, fatulence, colic, vomiting, diarrhoea, cough.(Basak et al,2017)
15	<i>Cassia fistula</i>	Indian laburnum	Fabaceae	The bark is used in treatment of inflammatory swellings and as a cleaning agent for ulcers and wounds. It is believed to decrease purulent discharge and act as local antiseptic. The seeds are are antibilious, asperitif, carminative and laxative (Ajaya Kumar <i>et al</i> ,2017)

RESULTS AND DISCUSSION

All the selected aqueous seed extracts exhibited various levels of antibacterial activity against *P.carotovorum*, the test bacterial species(Table.2) The aqueous seed extracts of *R.serpentina* exhibited highest length of inhibition zone .13.53 (Fig.1) followed by *A.subulatum* (11.33mm) against the test bacterial species .The inhibition zone was found to be10.0mm in seed extracts of *T.chebula* while 9.30mm in *F.vulgare* ,8.68 in *C.cyminum* and 8.03mm in *N.sativa*. There was no significant difference in

antibacterial activity between *A aromaticum* (7.97mm) and *A.indica* (7.92mm). The antibacterial activity was as par as in *T.bellirica* (7.00mm) and *C.viscosa* (7.01mm) while in *C. sativum* it was 7.28mm. The seed extracts of *P.nigrum*, *C.fistula* and *C.tora* expressed same activity (6.95 mm). Minimum inhibition zone (Fig.2) was observed in *E.officinallis* (6.68mm). No zone of inhibition of bacterial growth was observed in control (Fig.3).The inhibition zone in aqueous seed extracts ranged from 6.68mm to 13.53mm (Fig.4)




Fig. 4: Histogram showing zone of inhibition of *P.carotovorum* due to aqueous seed extracts of different plant species.

Table 2: Inhibition of growth of bacterial species by aqueous seed extract under *in vitro* condition.

Sl.no	Scientific Name	Common name (English)	Local name	Diameter in mm
1	<i>Terminalia chebula</i>	Chebulic myrobalan	Harida	10.00(3.24)
2	<i>T.myrobalan</i>	Beleric myrobalan	Bahada	7.00(2.74)
3	<i>Emblica officinals</i>	Aonla	Aonla	6.68(2.68)
4	<i>Amomum subulatum</i>	Greater cardamom	Bada alaicha	11.33(3.44)
5	<i>Rauvolfia serpentina</i>	Snake root	Patal garuda	13.63(3.72)
6	<i>Cleome viscosa</i>	Wild mustard	Banasorisha	7.01(2.74)
7	<i>Coriandrum satium</i>	Coriander	Dhania	7.28(2.79)
8	<i>Cuminum cyminum</i>	Cumin	Jeera	8.68(3.03)
9	<i>Azadirachata indica</i>	Neem	Nimba	7.92(2.86)
10	<i>Cassia tora</i>	Senna tora	Chhotachakunda	6.95(2.73)
11	<i>Nigella sativa</i>	Black cumin	Kalajeera	8.03(2.92)

12	<i>Foeniculum vulgare</i>	Fennel	Panamadhuri	9.30(3.13)
13	<i>Piper nigrum</i>	Black pepper	Golamaricha	6.95(2.73)
14	<i>Amomum aromaticum</i>	Aromatic cardamom	Alaicha	7.97(2.91)
15	<i>Cassia fistula</i>	Indian laburnum	Sunari	6.95(2.73)
16	Sterilized Water			0.00(0.71)
	SE(m) \pm			0.07
	CD(P=0.05)			0.21

Figures in parentheses are in $\sqrt{x+0.5}$ transformed values




Fig.1 : Inhibition zone observed in *R. serpentina*, Fig.2-Inhibition zone observed in *E.officinallis* ,Fig.3 – Control.

Several workers reported on the antimicrobial properties of above mentioned seed extracts and also the different parts of respective plants. Negi *et al* (2014) recorded the antibacterial activities of *R. serpentina*. In *A. subulatum* bio-chemical and biological activities had been studied by Bisht *et al*(2011). Antibacterial activity of black myrobalan (*Terminalia chebula*) against *Helicobacter pylori* had been studied and reported (Malekzadeh *et al* 2001). Rathre and Qureshi (2016) compiled the traditional uses and pharmacological behaviour of *F.vulgare*. Lacobellis *et al* 2005 reported the antibacterial activity of *C.cyminum*. Parihar *et al* (2012) detected the antioxidant immunomodulatory and antimicrobial activity of *Amomum aromaticum* against *Klebsiella pneumonia*. Dharmaratne *et al* (2018) reported antibacterial properties of *Tbellirica* against selected multi drug resistant bacteria. Datta and Kundabala, (2013) studied the antimicrobial efficacy of endodontic irrigants from *Azadirachta indica*. Saeed and Tariq (2007) reported the antimicrobial activities of *Emblca officinalis* and *Coriandrum sativum* against gram positive bacteria and *Candida albicans*. Kalane *et al* (2011) studied the antimicrobial activity of *Cassia tora*. Zou *et al*(2015) worked on antibacterial mechanism and activities of black pepper chloroform extract.

Antibacterial activity of black myrobalan (*Terminalia chebula*) against *Helicobacter pylori* had been studied and reported (Malekzadeh *et al* 2001). Mnif and Aifa (2015) compiled the beneficial effect of cumin(*Cuminum cyminum* L.) from traditional uses to potential

biomedical applications. Raja Ratna Reddy *et al*(2016) recorded the antimicrobial activity of *Azadirachta indica* (neem) leaf, bark and seed extracts.

The inhibition zone in aqueous seed extracts ranged from 6.68mm to 13.53mm (Fig.4).It was indicated all the seeds used in the test had antibacterial properties against *P. carotovorum*.The antibacterial property of *T. chhebula* (*Chhebolic myrobalan*), *T. belerica* (*Beleric myrobalan*), *E. officinalis* (Indian goose berry), (*A. subulatum*), (*Greater cardamom*), *R. serpentina* (Snake root), *C. sativum* (Dhania), *C. cyminum* (Cumin), *N. sativa* (Black cumin), *Foeniculum vulgare* (Fennel), *Piper nigrum* (Black pepper), *A. aromaticum* (aromatic cardamom) against *P. carotovorum* are reported to be new in India.Use of such extracts as tuber treatment at planting time, basal drenching in field and also tuber treatment in storage after harvest are to be tested.

REFERENCES

1. Ajay Kumar K, Satis S, Sayeed I, Hedge K Therapeutic uses of *Cassia fistula*:Review. *Int. J. Pharma. Chem. Res.*, 2017; 31(1): 38-43.
2. Ali BH,Blunden G Pharmacological and toxicological properties of *Nigella sativa*. *Phytother. Res.*, 2003; 17(4): 299-305.
3. Al-Timimi LN Antibacterial and anticancer activities of Fenugreek seed extract. *Asian Pacific J. Cancer Prevent*, 2019; 209(12): 3771-3776.

4. Ardha Jyoti,K. And Suba Rao,B. *In vitro* antibacterial activity of *Cleome viscosa*. *Pharma Sci.Monitor Repr*, 2010; 1(2): 71-78.
5. Basa SJ,Jayashankar Reddy V,Koshma M,Hanumanthu G,Dadakhalar S. A review on *Terminalia chebula*.*Inter. J. Pharmaco. Res*, 2017; 7(10): 187-191.
6. Basak, S.,Sharma,G.C.,Rangan, L. Ethnomedical uses of Zingiberaceous plants of North East India.*J. Ethonopharmaco*, 2010; 132(1): 286-296.
7. Bereksi MS,Hassaine H,Bekhechi C,Abdelouahid DE. Evaluation of antibacterial activity of some medicinal plant extracts commonly used in Algerian Traditional Medicine against some Pathogenic Bacteria. *Pharmacogon. J.*, 2018; 10(3): 507-512.
8. Bisht VK, Negi JS, Bhandari AK, Sunderiyal RC. *Ammomum subulatum* Roxb: Traditional biochemical and biological activities. *African J. Agricultu. Res.*, 2011; 6(24): 5386-5390.
9. Biswal G and Dhal, NK Survey on natural occurrence of diseases in potato at various locations of Odisha. *Internat. J. Plant Protect*, 2013; 6(1): 188-191.
10. Butt MS,Pasha I,Sultan MT,Randhwa MA,Saeed F,Ahmed W. Black pepper and health claims a comprehensive treatise. *Crit. Rev. Food. Sci.*, 2012; 53: 875-886
11. Kalane,V.J.and Chavan,R.T., Kalane V.J.and Deshmkuh,V.L. Antimicrobial activity of *Cassia tora*.*Curr.Bot.*, 2011; 2(1): 8-9.
12. Cowan MM Plant products as microbial agents. *Clinical Microbial Reviews*, 1999; 2(4): 63-78.
13. Dharmaratne MPJ,Manoraj A,Thevanesam V,Ekanayake A,Kumar NS,Liyanapathirana V,Abeyratne E and Bandara BR *Terminalia bellirica* fruit extracts : *in vitro* antibacterial activity against selected multidrug resistant bacteria,radical scavenge activity and cytotoxicity study on BHK-21 cells. *BMC Complementary and Alternative Medicine*, 2018; 18(1): 1-12.
14. Dutta A,Kundabala M Antimicrobial efficacy of endodontic irrigants from *Azadirachta indica*: An *in vitro* study. *Acta Odontologica Scandinavica*, 2013; 71(6): 1594-1598.
15. Fern K. a Tropical Plant Database,tropical.the ferns.info.2020-09-03< tropical. the ferns.info<view tropical.php, 2020.
16. Fern K. b Tropical Plant Database,tropical.the ferns.info.2020-09-04< tropical. The ferns.info<view tropical. php, 2020.
17. Gomez, K.A. and Gomez, A.A. Statistical procedure for Agricultural Research. John Wiley and Sons. New York, 1984; 680.
18. Kalane,V.J.and Chavan,R.T, Kalane ,V.J.and Deshmkuh, V.L. Antimicrobial activity of *Cassia tora*.*Curr.Bot.*, 2011; 2(1): 8-9.
19. Khan, KH. Role of *Embllica officinalis* in medicine-A review. *Botany Res. Internat*, 2009; 2(4): 218-228.
20. Kumar N and Khurana SM.Phytochemistry and medicinal potential of the *Terminalia bellirica*.*Indian J.Natural Products Resources*, 9(2): 97-107.
21. Lacobellis NS,Cantore PL,Capasso F, Senatore F. Antibacterial activity of *Cuminum cyminum* L and *Carum Carvi* L. Essential oils.*J.Agric.Food Chem*, 2005; 53(1): 57-61.
22. Mali RG A review on ethnobotany ,phytochemistry and pharmacology.*Pharmaceutical Biol.*, 2010; 48(1): 105-112.
23. Maity ,T.K.,Mandal,S.C.,Mulkherjee P, Saha., B.P., Das, J., Pal, M and Saha,B.P. Studies on anti-inflammatory effect of *Cassia* leaf extract.*Phytother.Res.*, 1998; 12(3): 221-223.
24. Malekzadeh,F,Ehsanifar,H,Shahamat,M,LevinM,ColwellR.R. Antibacterial activity of black myrobalan (*Terminalia chebula* against *Helicobacter pylori*.*Int. J.Antimicrob Agents*, 2001; 18: 85-88.
25. Mnif S and Aifa S. Cumin(*Cuminum cyminum* L.) from traditional uses to potential biomedical applications. *Chem. Biodivers*, 2015; 12(5): 733-42.
26. Negi JS,Bisht VK. Bhandaria AK, Bisht DS. Quantification of reserpine content and antibacterial activity of *Rauolfia serpentina* (L) Benth .ex Kurz.*African J. Microbiol. Res.*, 2014; 8(2): 162-166.
27. Newman DJ, Cragg GM and Snader KM The influence of natural products upon drug discovery; *Nat. Products Reports*, 2000; 17: 215-234.
28. Parihar L., Sharma,L., Kapoor,P., Parihar,P. Detection of antioxidant Immunomodulatory and antimicrobial activity of *Amomum aromaticum* against *Klebsiella pneumonia* *J.Pharm Res.*, 2012; 5(2): 901-905.
29. Perumal Samy,R,Ignacimuthu and Patric Raja,D. preliminary screening of ethnomedicinal plants from India.*J Ethnofarmacol.*, 1999; 66(2);: 235-240
30. Ponnusamy, K., Ramadevi, S.R. and Hopper W. Antibacterial activity of *Terminalia chebula* Retz.fruit extract. *African J. Microbiol. Res.*, 2009; 3(4): 180-184.
31. Raja Ratna Reddy Y, Krishna Kumari C,Lokanatha O,Mamatha S,Damodar Reddy C. Antimicrobial activity of *Azadirachta indica* (neem) leaf ,bark and seed extracts.*Int.J.Res.Phytochem.Phrmacol*, 2016; 3(1): 1-4.
32. Rathre MA,Qureshi M *Foeniculum vulgare*:A comprehensive review of its traditional use,phytochemistry, pharmacology and safety.*Arabian J. Chemistry*, 2016; 9(2): 51574-51583.
33. Saeed S and Tariq P. Antimicrobial activities of *Embllica officinalis* and *Coriandrum sativum* against gram positive bacteria and *Candida albicans*.*Pak J. Bot.*, 2007; 39(3): 913-917.
34. SharmaSK,Perianayagam BJ, Joseph A,Christina AJM. Antiinflammatory activity of ethanol an aqueous extracts of *Embllica officinalis* Gearth fruits *Hamdard Medicus*, 2003; 46: 75-78.

35. Somani AK and Shekhawat GS Bacterial soft rot of potato in India. Technical Bulletin. 21. CPRI, Shimla, 1990; 32.
36. Waheed, A., Miana G.A.Ahmad SI,Khan MA. Clinical investigation of hypoglycemic effect of *Coriandrum sativum* in type-2 (NIDDM) diabetic patients. *Pakistan J. Pharmacol*, 2006; 23(1): 7-11
37. Valgas C, Souza SM, Smania EFA,Smania A. Screening methods to determine antibacterial activity of natural products. *Brazilian J. Microbiol*, 2007; 38(2): 369-380.
38. Venugopalan SK and Visweswaran N. Neem (*Azadirachta indica*): Prehistory to contemporary medicinal uses to humankind. *Asian Pac J Trop Biomed*, 2013; 3(7): 505-514.
39. Zou L, Hu Y-Y, Chen W-X. Antibacterial mechanism and activities of black pepper chloform extract. *J Food Sci Technol.*, 2015; 52(12): 8196-8203.