

**PRE-OPERATIVE PREDICTION OF DIFFICULT LAPAROSCOPIC
CHOLECYSTECTOMY BY ESTIMATION OF HIGH SENSITIVE C - REACTIVE
PROTEIN (HS-CRP) ALONG WITH ULTRASONOGRAPHIC FINDINGS- A STUDY****Dr. Abhijit Sarma and *Dr. Purujit Choudhury**

Associate Professor of Surgery, Gauhati Medical College.

Received on: 09/10/2021

Revised on: 29/11/2021

Accepted on: 19/12/2021

*Corresponding Author

Dr. Purujit ChoudhuryAssociate Professor of
Surgery, Gauhati Medical
College.**ABSTRACT**

Background: Laparoscopic cholecystectomy is considered to be the gold standard for the treatment of gallstones. However, conversion of laparoscopic cholecystectomy to open cholecystectomy is done by surgeons in difficult cases. The aim of the study is to use Hs-CRP level in correlation with ultra-sonographic findings to predict difficult laparoscopic cholecystectomy pre-operatively. It would help in anticipating a difficult laparoscopic cholecystectomy preoperatively, better counselling of patients, avoiding unnecessary laparoscopic dissection and hence less morbidity. **Materials and Methods:** It is an original study conducted in the department of general surgery of Gauhati Medical College and Hospital. The study group included 40 consecutive patients undergoing laparoscopic cholecystectomy for cholelithiasis. Cholelithiasis was diagnosed on the basis of ultrasonography done preoperatively. Hs-CRP level was determined pre-operatively and postoperatively after 6hrs and 12hrs. It was correlated with ultra-sonographic findings to predict difficult cholecystectomy. **Results:** The mean value of Hs-CRP was found to be 9.68 mg/L preoperatively, 21.88 mg/L at 6 hours and 26.57 mg/L at 12 hours. While the mean value of Hs-CRP was found to be significantly higher in cases which needed conversion. USG findings such as presence of adhesion, GB wall thickness more than 3mm, and narrow Calot's triangle, were found to be significant predictors for conversion. On correlation, the positive predictive value increased. **Conclusion:** Pre-operative values of Hs-CRP when combined with ultrasonography findings, can predict difficult cholecystectomy. It will help in better selection of patients, and help in decreasing morbidity.

KEYWORDS: Laparoscopic cholecystectomy, Hs-CRP, Ultrasonography, Conversion.

BACKGROUND

Gallstones are among the most common gastrointestinal illness requiring surgery. They are a major cause of morbidity and mortality throughout the world.^[1] Laparoscopic cholecystectomy has been recognized as the gold standard for the treatment of gallstones.^[2] Currently it is estimated that over 75% of cholecystectomies are performed using laparoscopic approach.^[3]

The advantages include earlier return of bowel function, less postoperative pain, improved cosmesis, shorter length of hospital stay, earlier return to full activity and decreased overall cost.^[3,4,5] However very often surgeons face a number of surgical and technical difficulties while performing the procedure due to which the procedure has to be converted into open cholecystectomy. The conversion rate for elective laparoscopy cholecystectomy varies from 2%-15%.^[6,7,8,9] Inability to define the anatomy and difficult dissection are the leading reasons for conversion followed by other complications like

bleeding. It has now become one of the most common operation performed by general surgeons.^[10]

CRP is an acute phase reactant protein synthesized by the liver in response to factors released by macrophages and adipocytes in the face of inflammation. During the acute phase response, level of CRP rapidly increases within 2hrs of acute insult, reaching a peak at 48hrs. CRP declines with a relatively short life of 19hrs. The Hs-CRP assay is an advanced quantitative analysis using laser nephelometry. It can measure very low levels of C reactive proteins. It has been shown to be more accurate in predicting risk of morbidity in undertaking lap cholecystectomy in normal people than standard CRP.

Till today, ultrasound has been a widely used modality for predicting the difficult laparoscopic cholecystectomy in pre-operative patients. Due to its several advantages like safety, wide availability, etc it is preferred as an initial imaging study. Ultrasound shows stones in the gallbladder with a sensitivity and specificity of more than 90%.

This study was undertaken to see if there is any correlation of Hs-CRP in assisting and improving the accuracy of ultrasonography in predicting difficult Laparoscopy. This would help a surgeon to anticipate better a difficult cholecystectomy, better pre-operative counselling of patients, avoid excessive intraoperative manipulations, go for conversions early, reduce time of surgery and hence also minimize post-operative complications.

MATERIALS AND METHODS

This was an original study done in Gauhati medical college and hospital in Guwahati, Assam. The study period was from 1/7/18 to 30/6/19. The study population comprised of 40 consecutive patients undergoing laparoscopic cholecystectomy after fulfilling the inclusion criteria. Patients having Asymptomatic gallstones, acalculous cholecystitis, associated choledocholithiasis, malignancy, pregnancy, patients under steroid therapy/local tissue irradiation, coagulopathies, previous operation of hepatobiliary system or upper GI surgery, any associated illness contraindicating cholecystectomy surgery and any associated illness leading to rise in CRP levels, were excluded from the study. Informed consent was taken. Patients were also informed that in case of much difficulty, laparoscopy would be converted to open cholecystectomy. Cases were diagnosed as having cholecystitis due to cholelithiasis on the basis of ultrasonographic findings. The cases were prepared for laparoscopic cholecystectomy as per as protocol followed in our institution. Relevant investigations were done. It included random blood sugar, complete hemogram, Liver function test, renal function test, thyroid function test, chest x-ray, ECG, viral markers and ultrasonography of whole abdomen. The Hs-CRP level was determined pre-operatively and 6hrs and 12hrs. 5ml venous blood was collected in clot vials for transfer to biochemistry laboratory. The samples were analyzed by the help of kit made by ROSS co., Germany for COBAS INTEGRA 400 analyzer. At the time of admission, a thorough history, general and clinical examination was carried out and records were maintained. The age group was between 21-60yrs.

Surgical Procedures

All the surgeries were performed by consultant and done under general anesthesia. Pneumoperitoneum is first created by using a Verres needle through an infraumbilical port. A total of 4 ports were used. 2 of 10mm size and 2 of 5mm size. Adhesion was released first. Dissection was done to make the Calot's triangle naked. The cystic artery and cystic duct was separately clipped and divided. The gallbladder was the dissected off the gallbladder fossa, using the monopolar cautery hook. At the completion of surgery, gallbladder was taken out using the epigastric port. The port was enlarged if needed. A saline lavage was done. Hemostasis was achieved. Drain was placed whenever clinically

indicated. Ports were closed. When conversion was needed, it was done through right subcoastal incision.

Statistical Analysis

Normally distributed continuous variables were compared using the paired t- test. Categorical variables were analysed using the Fischer exact test. For all statistical test, a p-value less than .05 was taken as significant. ROC analysis curve was plotted for values of Hs-CRP and its sensitivity for prediction of conversion.

RESULTS

Patients Demographic: The total number of cases were 40 with 11 males and 29 females. Gender and age wise these were comparable as the p value was found to be 1.000 using Fischer exact test. The male: female ratio was found to be 1:2.6. there were total of 8 cases in the age group of 21-30yrs, 13 cases were in the age group of 31-40yrs, 16 in the age group of 41-50yrs and 3 in 51-60 yrs. The mean age of incidence was found to be 39.65 yrs.

Conversion to open Cholecystectomy

Out of 40 cases, 3 cases required conversion to open cholecystectomy. Among these 3 cases, 1 is male and 2 are females, p value was found to be 1.000, which meant the difference is insignificant. The cases which required conversion were equally divided across the age group of 31-40, 41-50, 51-60 yrs. The conversion in respect to age group and gender was found to be insignificant. However, to comment definitively we need a larger sample size.

Conversion in Relation to USG

For a thickened gallbladder, the cut-off value was taken to be 3mm radiologically. Thickened gallbladder was found in 7 cases. Out of which 3 cases required conversion to open cholecystectomy. The p-value was found to be 0.0035 which reflects GB wall thickness as a significant risk factor. On USG, 6 cases had adhesion present. Out of which 3 cases needed conversion to open cholecystectomy. The presence of adhesion is a significant risk factor for conversion. In 5 cases, Calot's triangle was observed to be narrowed radiologically, out of which 3 cases required conversion. The p value was found to be 0.0010. It reflects narrow Calot's triangle as a significant risk factor for conversion. The p value of the difference in conversion rate among patients with single gallstones was found to be 0.498 using Fischer exact test. Therefore, the difference in conversion rate among patients having single or multiple calculi is not significant. In relation to number of calculi in gallbladder seen radiologically it was found that one case had single calculi and two cases had multiple calculi, among the three cases which required conversion. However, the p value of the difference in conversion rate among patients with single gallstone and multiple gallstone was found to be 0.498 using Fischer exact test, which is not significant.

Conversion In Relation To Hs-Crp

The mean pre-operative value of Hs-CRP was found to be 9.68mg/l. It increased to 21.88 mg/l 6hrs post-operatively and increased to 26.57 mg/l 12 hrs postoperatively. The mean pre-operative value of Hs-CRP in cases which needed conversion was found to be 65.3mg/L pre-operatively, 118.9mg/L 6hrs post-operatively, 129mg/L 12 hrs post-operatively. The difference in values of Hs-CRP between the cases who underwent laparoscopic cholecystectomy successfully and converted cases were found to be significant. The presence of high level of Hs-CRP gives a very good indication of ongoing inflammation in the body. Studies have shown that in case of inflammation there is rise in level of CRP within 4-6 hours. It reaches a peak value in two days and thereafter declines. Therefore, the values of Hs-CRP are also significant in understanding recovery as it should decline after 2 to 3 days in absence of any infection postoperatively.

On using a cut off value of Hs-CRP more than 20mg/l preoperatively, sensitivity was found to 100% and specificity was 94.5%. As the cut off value was increased progressively upto 50mg/l, the sensitivity fell to 66.66% and specificity remained same at 100%.

On combining USG parameters with different Hs-CRP cut off values, it is observed, as we increase the cut off from 20mg/l to 50mg/l, the sensitivity remains same for 20mg/l, 30mg/l, and 40mg/dl. However it decreases for 50mg/l. Both positive and negative predictive values were serially analysed with different cut off points for Hs-CRP values. An improvement in positive predictive

value is observed, as the cut-off of Hs-CRP concentration is raised while negative predictive value remained at 100 percent and fell to 97.36 percent for cut-off value of 50 mg/dl. It is done between these two parameters only, as this was aim of the study.




Fig 1: Age and gender distribution of study group.




Fig 2: Showing the specificity-sensitivity analysis for conversion from LC to OC using different cut off of Hs - CRP from more than 20 mg/l through 30mg/l, 40mg/l, 50mg/l.

Table 1: Mean and standard deviation of values of Hs- CRP, preoperatively and post operatively at 6 Hrs and 12 Hrs.

Group parameter	N	mean	Standard deviation
Cases- pre op CRP	40	9.68	±17.66
Cases- post op at 6 hrs CRP	40	21.88	±31.2
Cases- post op at 12 hrs CRP	40	26.57	±32.7

Table 2: Sensitivity, specificity, PPV and NPV of ultrasonography combined with different pre- op cut off values of hs-CRP.

Parameters	Positive findings on USG and hs-crp>20mg/l	Positive findings on USG And hs-CRP>30mg/l	Positive findings on USG and hs-CRP>40mg/l	Positive findings on USG and hs-CRP >50 mg/l
Sensitivity (%)	100%	100%	100%	66.6%
Specificity (%)	98.7%	99.3%	100%	100%
Positive predictive value (%)	17.6%	18.75%	20%	21.4%
Negative predictive value (%)	58.7%	57.8%	56.9%	56.06%


Fig. 3: ROC CURVE.

Case Processing Summary

VAR00001	Valid N (listwise)
Positive ^a	3
Negative	37

Larger values of the test result variable(s) indicate stronger evidence for a positive actual state.

a. The positive actual state is 1.00.



Area Under the Curve

Test Result Variable(s):VAR00002		Asymptotic 95% Confidence Interval		
Area	Std. Error ^a	Asymptotic Sig. ^b	Lower Bound	Upper Bound
1.000	.000	.004	1.000	1.000
a. Under the nonparametric assumption				
b. Null hypothesis: true area = 0.5				

Coordinates of the Curve

Test Result Variable(s):VAR00002

Positive if Greater Than or Equal To ^a	Sensitivity	1 - Specificity
.2000	1.000	1.000
1.2500	1.000	.919
1.3650	1.000	.892
1.5150	1.000	.865
1.6500	1.000	.838
1.7500	1.000	.811
1.8500	1.000	.784
1.9500	1.000	.730
2.1000	1.000	.703
2.2500	1.000	.676
2.3500	1.000	.622
2.5300	1.000	.595
2.7300	1.000	.568
2.9500	1.000	.514
3.1500	1.000	.432
3.2500	1.000	.405
3.3500	1.000	.378
3.5000	1.000	.351
3.9500	1.000	.324
4.6500	1.000	.297
5.4500	1.000	.270
6.1500	1.000	.243
6.7500	1.000	.216
7.1500	1.000	.189
7.8500	1.000	.162

9.2000	1.000	.135
10.5000	1.000	.108
12.7500	1.000	.081
18.5000	1.000	.054
27.1000	1.000	.027
39.8000	1.000	.000
54.8500	.667	.000
73.9500	.333	.000
87.2000	.000	.000

a. The smallest cutoff value is the minimum observed test value minus 1, and the largest cutoff value is the maximum observed test value plus 1. All the other cutoff values are the averages of two consecutive ordered observed test values.

DISCUSSION

The male: female ratio of occurrence of gallstones was found to be almost equal to 1:3. This is in agreement with studies done by NovacekG,^[11] where occurrence of gallstones was found to be 2 to 3 times higher. Moghaddam AA et al^[12] reported a 2.73 times higher incidence of gallstones in female. The conversion rate in our study was found to be 7.5%. this is comparable to a conversion rate of 1.8% - 27.7% as reported by different series. Kanann SA et al^[13] reported a conversion rate of 10% in acute cholecystitis and 4% in chronic cholecystitis. In our study, there was no relation found between gender and age of patient with conversion. This is in contrast to several other studies. Shapiro AJ, Lein HH, and Huang CS.^[14,15] where male gender is found to be a risk factor. Age has been recognized as an independent risk factor for conversion in studies done by Liu CV,^[3] Kama NA,^[16] S Ibrahim,^[17] et al. However number of cases is a limitation in our study to make a definitive comment regarding age and sex with risk of conversion. Morbid obesity and presence of diabetes mellitus has been shown to be risk factors for conversion in studies of S Ibrahim et al and lein HH. But in our study, these factors were not included as these patients were excluded. Gallbladder wall thickness more than 3 mm on ultrasonography was found to be a predictor for conversion. It is in agreement with studies done by U Jethwani, Shapiro AJ, and Tayeb M.^[14,18,19] Other findings such as adhesion and narrow Calot's triangle were found to be significant predictor for conversion. This is in accordance to other studies done by Volkan G (2011), Tayeb et al, Shapiro et al, Chand P et al.^[14,19,20,21] Levels of Hs-CRP were determined preoperatively and post operatively at 6hrs and 12 hrs. Our study concludes that high Hs-CRP values preoperatively predict conversion. The ROC curve reveals the highest sensitivity for Hs-CRP to be around 40mg/l for conversion. This is in agreement with other studies done, like Schafer M et al, Kam Wa et al, Weavers KP et al, and Singh BA et al,^[22,23,24,25] where high values of CRP was found to be significant predictors for conversion.

CONCLUSIONS

The study was successful in establishing that, when the pre-operative values of Hs-CRP is correlated with ultrasonography findings, it can predict difficult cholecystectomy. It will help surgeons in making better judgement, and better counselling of patients prior to surgery keeping co-morbid conditions in view. Hs-CRP with ultrasound findings and experience of the surgeon will help in better selection of patients, thus reducing morbidity and mortality.

BIBLIOGRAPHY

1. Becker H, Buess G. The European experience with laparoscopic cholecystectomy. *Am J Surg*, 1991; 161(3): 385-7.
2. Alponat A, Kum CK, Koh BC. Predictive factors for conversion of laparoscopic cholecystectomy. *World J Surg*, 1997; 21: 629-33.
3. David EJ, Marshall MK. Pathogenesis and treatment of gallstones. *N Engl J Med*, 1993; 328: 412-21.
4. Brunicardi FC. Gallbladder and extrahepatic biliary system. In: Corbett S, editor. *Schwartz Principles of Surgery*, 10 edn. United States of America: Mc Graw Hill, 2015; 1319.
5. Demestral C, Roststein D, Laupacis A, Hoch JS, Zagoski B, Alali AS et al. Comparative operative outcomes of early and delayed cholecystectomy for acute cholecystitis ; a population based propensity score analysis. *Ann Surg*, 2014; 259: 10-5.
6. Wu XS, Shi LB, Gu J, Dong P, Lu JH, *et al.* Single incision laparoscopic cholecystectomy versus multi-incision laparoscopic cholecystectomy: A meta-analysis of randomized clinical trials. *J Laparoendosc adv surg tech*, 2013; 23: 183-91.
7. Cuschieri A, Dubois F, Mouiel P, Mouret P, Becker H, Buess G *et al.* The European experience with laparoscopic surgery. *Am J Surg*, 1991Mar; 161(3): 385-7.
8. Wiebke EA, Pruitt AI, Howard TJ. Conversion of laparoscopic to open cholecystectomy. An analysis of risk factors. *SurgEndosc*, 1996; 10: 742-5.
9. Michael R, Fred B, Ponsky J. predictive for conversion of laparoscopic cholecystectomy. *Am J Surg*, 2002; 184(3): 254-8.
10. Giger UF, Risk factor for perioperative complications in patients undergoing laparoscopic

- cholecystectomy: analysis of 22,953 consecutive cases from the Swiss Association of Laparoscopic and Thoracoscopic Surgery Database. *J Am Coll Surg*, 2006; 203(5): 723-8.
11. Das S, Biliary system. In: Das S, editor. A concise textbook of surgery. Edn. 6th Kolkata: S.R Das, 2010.
 12. Novacek G. Gender and gallstones. *Wien Med Wochenschr*, 2006; 156(19-20): 527-33.
 13. Moghaddan AA, Khoram A, Bonjar MM, Mohammadi M, Ansari H. The prevalence and risk factors of gallstones among adults in south-east of Iran: A population based study *Glob J health sci.*, 2016Apr; 8(4): 60-7.
 14. Kanaan SA, Murayama KM, Merriam LT, Daves LG, Prytowsky JB, Rege RV *et al.* Risk factor for conversion of laparoscopic cholecystectomy to open cholecystectomy. *J Surg Res.* Elsevier science, 2002.
 15. Shapiro AJ, Costello C, North JH. Predicting conversion of laparoscopic cholecystectomy for acute cholecystitis. *J soc laparoendosc. Cholecystectomy for acute cholecystitis*, 1999Apr-Jun; 3(2): 127-30.
 16. Lein HH, Huang CS. Male gender: risk factor for severe symptomatic cholelithiasis. *World J Surg*, 2002 may; 26(5): 598-601.
 17. Kama NA, Kologlu M, Doganay M, Reis E, Athi M, Dolapci MA. A risk score for conversion from laparoscopic to open cholecystectomy. *Am J Surg*, 2001; 181: 520-5.
 18. Ibrahim S, Hean TK, Hol S. Risk factor for conversion to open surgery in patients undergoing laparoscopic cholecystectomy. *World J Surg*, 2006 Sep; 30(9): 1698-704.
 19. Jethwani U, Singh G, Mohil RS, Kandwal V, Razdan S, Chouhan J *et al.* Prediction of difficulty and conversion in laparoscopic cholecystectomy. *OA Minim Invasive Surg*, 2013Aug; 1(1): 2.
 20. Tayeb M, Raza SA, Khan MR, Azami R. Conversion from laparoscopic cholecystectomy to open cholecystectomy. Multivariate analysis of pre-op risk factors. *J Postgrad Med*, 2005; 51(1): 17-20.
 21. Volkan G, Sulaimanov M, Harzinedaroglu SM. What necessitates the conversion to open cholecystectomy? A retrospective analysis of 5164 consecutive laparoscopic operation. *Clinics*, 2011Mar; 66(3): 417-20.
 22. Chand P, Singh R, Yadav M, Singla RL, Singh B. Pre-operative USG as a predictor of difficult laparoscopic cholecystectomy that requires conversion to open procedure. *Niger J Surg*, 2015Jul-dec; 21(2): 102-5.
 23. Schafer M, Krahenbuhl L, Buchler MW. Predictive factors for the type of surgery in acute cholecystitis. *Am J Surg*, 2001; 182: 291-7.
 24. Wevers KP, Van Westren HI, Patijn GA. Laparoscopic cholecystectomy in acute cholecystitis: C-RP level combined with age predicts conversion Surg Laparosc Endosc Percutan Tech, 2013Apr; 23(2): 163-6.
 25. Wa K, Mok J, Goh Y G, Ravindra S. Is C reactive protein the single most useful predictor of difficult laparoscopic cholecystectomy or its conversion? A pilot study. *J Minim Access Surg*, 2016Jan-Mar; 12(1): 26-32.
 26. Arora BS, Sen P, Singh RK, Biswal I, Paruthy SB. Hs-CRP levels as predictor of difficult laparoscopic cholecystectomy- current status evaluation experience in a teaching institution. *Int Surg J*, 2017; 4(4).